

A LOGARITHMIC BOUND FOR THE CHROMATIC NUMBER OF THE ASSOCIAHEDRON

LOUIGI ADDARIO-BERRY, BRUCE REED, ALEX SCOTT, AND DAVID R. WOOD

Abstract. We show that the chromatic number of the n -dimensional associahedron grows at most logarithmically with n , improving a bound from and proving a conjecture of Fabila-Monroy et al. [*Discrete Math. Theor. Comput. Sci.*, 2009].

1. Introduction

The associahedron \mathcal{A}_n is an $(n - 3)$ -dimensional convex polytope that arises in numerous branches of mathematics, including algebraic combinatorics [3, 5, 8, 13] and discrete geometry [1, 9, 10]. Associahedra are also called Stasheff polytopes after the work of Stasheff [13], following earlier work by Tamari [14]. We are only interested in the 1-skeleton of the associahedron, so we consider it as a graph, defined as follows.

The elements of the associahedron \mathcal{A}_n are triangulations T of the convex n -gon with vertices labeled by $\{0, \dots, n - 1\}$ in clockwise order. For any such a triangulation T , we always denote triangles of T by the sequence ABC of their vertices, ordered so that $A < B < C$. We write $E(T)$ for the set of edges contained in T . Every triangulation $T \in \mathcal{A}_n$ contains the edges $01, 12, \dots, (n - 1)0$; we refer to these as *boundary edges*. For $T \in \mathcal{A}_n$, each non-boundary edge $e \in E(T)$ is contained in a unique quadrilateral $Q = Q_T(e) = ABCD$ with $A < B < C < D$; we will always list the vertices of quadrilaterals in increasing order. *Flipping* the edge e means replacing e by the other diagonal of Q . (See Figures 1a and 1b.) Two triangulations $T, T' \in \mathcal{A}_n$ are adjacent in \mathcal{A}_n if they may be obtained from one another by a single flip.




Figure 1. Portions of two adjacent triangulations of an n -gon.

Graph-theoretic properties of associahedra have been well-studied. For example, it is easily seen that \mathcal{A}_n is $(n - 3)$ -regular. Hurtado and Noy [6] proved that \mathcal{A}_n is Hamiltonian and has connectivity $n - 3$, as well as determining its automorphism group. The diameter of \mathcal{A}_n and several

Date: November 21, 2018.

1991 *Mathematics Subject Classification.* 52B05, 05C15.

Louigi Addario-Berry is supported by an NSERC Discovery Grant and an FRQNT Team Research Grant.

Bruce Reed is supported by NSERC and the ERATO Kawarabayashi Large Graph Project.

Alex Scott is supported by a Leverhulme Trust Research Fellowship.

equivalent questions has been studied extensively [2, 11, 12]. Sleator et al. [12] proved that the diameter equals $2n - 8$ for sufficiently large n , and recently Pournin [11] showed that $2n - 8$ is the answer for $n \geq 7$.

This paper studies the chromatic number of \mathcal{A}_n , a quantity which was first considered by Fabila-Monroy et al. [4]. That work gave an explicit $\lceil \frac{n}{2} \rceil$ -colouring of \mathcal{A}_n , and observed that $\chi(\mathcal{A}_n) \leq O(n/\log n)$, based on the result of Johansson [7] which says that every triangle-free graph with maximum degree Δ is $O(\Delta/\log \Delta)$ -colourable. No non-constant lower bound on $\chi(\mathcal{A}_n)$ is known. Indeed, the best known lower bound is $\chi(\mathcal{A}_{10}) \geq 4$ [private communication, Ruy Fabila-Monroy]. Fabila-Monroy et al. [4] conjectured a $O(\log n)$ upper bound. We prove this conjecture.

Theorem 1. $\chi(\mathcal{A}_n) = O(\log n)$.

2. The Proof

We prove Theorem 1 by tracking how several carefully chosen properties of triangulations change when an edge is flipped. To see how this yields a route to bounding the chromatic number of \mathcal{A}_n , first recall that if f is a graph homomorphism from \mathcal{A}_n to some graph G , which is to say that $f : \mathcal{A}_n \rightarrow V(G)$ is adjacency-preserving, then $\chi(\mathcal{A}_n) \leq \chi(G)$. This fact may be generalized as follows. Suppose that $(G_i)_{i \in I}$ is a finite set of graphs and $(f_i : \mathcal{A}_n \rightarrow V(G_i))_{i \in I}$ are functions such that if $T, T' \in \mathcal{A}_n$ are adjacent triangulations then there exists $i \in I$ for which $f_i(T)$ and $f_i(T')$ are adjacent in G_i . For each $i \in I$, let κ_i be a proper colouring of G_i with $\chi(G_i)$ colours, and colour $T \in \mathcal{A}_n$ with the vector $(\kappa_i(f_i(T)))_{i \in I}$. If T and T' are adjacent then $(\kappa_i(f_i(T)))_{i \in I}$ and $(\kappa_i(f_i(T')))_{i \in I}$ differ in at least one coordinate; thus this is a proper colouring of \mathcal{A}_n , and $\chi(\mathcal{A}_n) \leq \prod_{i \in I} \chi(G_i)$. The remainder of the paper is devoted to defining the functions we will use, and showing they have the requisite properties.

Two fundamental notions that we will use are the *type* of a quadrilateral and the *scale* of an edge. For a quadrilateral $Q = Q_T(e) = ABCD$ contained within triangulation T , we say Q is *type-1* if $e = AC$, and otherwise say Q is *type-2*; we say that an edge e is *type-1* or *type-2* according to the type of the quadrilateral $Q_T(e)$. (For example, in Figure 1, $Q_T(AC) = ABCD$ is type-1 and $Q_T(BC) = ABEC$ is type-2, and in Figure 1, $Q_{T'}(BC)$ is type-1 and $Q_{T'}(BD)$ is type-2.)

We fix a parameter $r > 0$ to be chosen later (in fact we will end up taking $r = 3$). For an edge $e = UV$, define the *scale* of e to be $\sigma_e = \lceil \log_r |U - V| \rceil \in \{0, 1, \dots, \lceil \log_r(n - 1) \rceil\}$. The scales of the edges incident to triangles within a fixed quadrilateral Q will be a key input to the functions we define.

We first consider the effect of edge flips on triangles ABC where two of the three incident edges have the same scale. If AB (resp. BC , AC) is the unique edge whose scale is different from the others, then we say ABC is a *type- l* (resp. *type- m* , *type- r*) triangle. If all three edges have the same scale s , then we say ABC is a *type- z* triangle. Let (l_T, m_T, r_T, z_T) be the vector counting the number of type- l , type- m , type- r and type- z triangles in T .

For the remainder of the paper, we fix a triangulation T , and consider the effect of flipping a type-1 edge $e = AC$ within its quadrilateral $Q_T(e) = ABCD$, to form T' .

Proposition 2. *Suppose that $r \geq 3$, and that σ_{AC} , σ_{BD} and σ_{BC} are all different. Then $(l_{T'}, m_{T'}, r_{T'}, z_{T'}) \neq (l_T, m_T, r_T, z_T)$.*

Proof. By assumption, $\sigma_{BD} \neq \sigma_{AC}$ and $\sigma_{BC} < \min(\sigma_{AC}, \sigma_{BD})$. We argue by contradiction. To this end, suppose that $(l_{T'}, m_{T'}, r_{T'}, z_{T'}) = (l_T, m_T, r_T, z_T)$. Note that

$$\log_r(D - A) \leq \log_r(3 \max(B - A, C - B, D - C)) \leq 1 + \log_r \max(B - A, C - B, D - C).$$

Taking ceilings, it follows that

$$\sigma_{AD} \leq 1 + \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD}). \quad (1)$$

The preceding equation requires one of three inequalities to hold; the next three paragraphs treat the possibilities one at a time.

Suppose that $\sigma_{AB} = \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD})$. Then either $\sigma_{AC} = \sigma_{AB}$ or (using (1) and the fact that $\sigma_{AB} \leq \sigma_{AC} \leq \sigma_{AD}$) we have $\sigma_{AC} = \sigma_{AB} + 1 = \sigma_{AD}$.

If $\sigma_{AC} = \sigma_{AB}$, as in Figure 2a, then ABC is a type- m triangle so, since we assume the triangle type vector is unchanged by flipping edge AC , either ABD or BCD is also type- m . BCD is not type- m , as $\sigma_{BC} < \sigma_{BD}$ by assumption, so ABD is type- m and hence $\sigma_{AB} = \sigma_{AD}$. But then ABC and ACD are both type- m , which gives a contradiction as BCD is not.

If $\sigma_{AC} = \sigma_{AB} + 1$, as in Figure 2b, then ACD is type- m or type- z , so either ABD or BCD is type- m or type- z . But BCD is neither, as $\sigma_{BC} < \sigma_{BD}$, and ABD is neither as $\sigma_{AB} \neq \sigma_{AD}$.

Next suppose that $\sigma_{BC} = \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD})$, as in Figure 2c. Since $\sigma_{BC} \neq \sigma_{AC}$ and $\sigma_{BC} \neq \sigma_{BD}$ by assumption, and all scales are at most σ_{AD} , it must be that $\sigma_{AC} = \sigma_{BD} = \sigma_{AD}$; but this is ruled out by assumption.

Finally, suppose that $\sigma_{CD} = \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD})$. This case is the same as the first case, as we can apply the argument to a reversed copy of the associahedron (which exchanges type- l and type- m triangles, while leaving the other two types invariant). \square




Figure 2. Writing $\sigma = \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD})$, the subfigures correspond to possible configurations arising in the proof of Proposition 2.

Proposition 3. *If $\sigma_{AC} = \sigma_{BD} = \sigma_{BC}$ and $(l_{T'}, m_{T'}, r_{T'}, z_{T'}) = (l_T, m_T, r_T, z_T)$, then either $\sigma_{AD} = \sigma_{BC}$ or $\sigma_{AB} = \sigma_{BC} = \sigma_{CD}$.*

Proof. In this case ABC is type- l or type- z , and BCD is type- m or type- z . If ABC is type- l then since the triangle type vector doesn't change it must be that ABD is type- l , which implies that $\sigma_{AD} = \sigma_{BD}$, yielding the result in this case. Similarly, if BCD is type- m then it must be that ACD is type- m , which implies that $\sigma_{AD} = \sigma_{AC}$. Otherwise, both ABC and BCD are type- z , in which case we indeed have $\sigma_{AB} = \sigma_{BC} = \sigma_{CD}$. \square

For a triangulation $T \in \mathcal{A}_n$ and $k \in \{1, 2\}$ and $i \in \{0, 1, \dots, \lceil \log_r(n-1) \rceil\}$, let

$$s_i^k(T) = \#\{e \in E(T) : Q(e) \text{ is type-}k, \sigma_e = i\}.$$

We assign an integer label $c(T)$ to T given by

$$c(T) = \left(\sum_{i=0}^{\lceil \log_r(n-1) \rceil} 2is_i^1(T) + \sum_{i=0}^{\lceil \log_r(n-1) \rceil} 3is_i^2(T) \right) \pmod{3\lceil \log_r n \rceil}.$$

The utility of such a labelling rule is explained by the following fact. We continue to work with triangulations T and T' related by an edge flip within quadrilateral $ABCD$ with $AC \in E(T)$ and $BD \in E(T')$, as above.

Proposition 4. *If exactly two of σ_{AC} , σ_{BD} and σ_{BC} are equal then $c(T') \neq c(T)$.*

Proof. First suppose that BC is not a boundary edge, and let V be the unique vertex of T with $B < V < C$ which is incident to both B and C . Note that $ABCD = Q_T(AC)$ is type-1 in T and $ABCD = Q_{T'}(BD)$ is type-2 in T' . Also, $Q_T(BC) = ABVC$ is type-2 in T and $Q_{T'}(BC) = BVCD$ is type-1 in T' . It is not hard to check that no other quadrilaterals change type when moving from T to T' . We thus have

$$\begin{aligned} c_1(T') - c_1(T) &= 3\sigma_{BD} - 2\sigma_{AC} + 2\sigma_{BC} - 3\sigma_{BC} && \text{mod } (3\lceil \log_r n \rceil) \\ &= 3\sigma_{BD} - 2\sigma_{AC} - \sigma_{BC} && \text{mod } (3\lceil \log_r n \rceil). \end{aligned}$$

It follows that if $\sigma_{BC} < \sigma_{BD} = \sigma_{AC}$ then

$$c_1(T') - c_1(T) = \sigma_{BD} - \sigma_{BC} \pmod{3\lceil \log_r n \rceil} \neq 0;$$

the difference is non-zero mod $(3\lceil \log_r n \rceil)$ since all scales are at most $\lceil \log_r n \rceil$. Similarly, if $\sigma_{BC} = \sigma_{BD} < \sigma_{AC}$ then

$$c_1(T') - c_1(T) = 2(\sigma_{AC} - \sigma_{BD}) \pmod{3\lceil \log_r n \rceil} \neq 0,$$

and if $\sigma_{AC} = \sigma_{BC} < \sigma_{BD}$ then

$$c_1(T') - c_1(T) = 3(\sigma_{BD} - \sigma_{AC}) \pmod{3\lceil \log_r n \rceil} \neq 0,$$

Since $\sigma_{BC} \leq \min(\sigma_{AC}, \sigma_{BD})$, these are the only possibilities.

The case when BC is a boundary edge is very similar, but easier. In this case we obtain that

$$c_1(T') - c_1(T) = 3\sigma_{BD} - 2\sigma_{AC} \pmod{3\lceil \log_r n \rceil}.$$

Since BC is a boundary edge, AC and BD are not, so $\sigma_{BC} = 0$ and $\sigma_{AC} \neq 0$, $\sigma_{BD} \neq 0$. It follows by assumption that $\sigma_{AC} = \sigma_{BD}$, so

$$c_1(T') - c_1(T) = \sigma_{AC} \pmod{3\lceil \log_r n \rceil} \neq 0. \quad \square$$

Propositions 2, 3 and 4 imply that the label $c(T)$ and the type vector (l_T, m_T, r_T, z_T) together distinguish T from T' except in the following cases.

- (1) AC, BD, BC , and AD have the same scale and AB, CD have smaller scales.
- (2) AC, BD, BC, AD and AB have the same scale and CD has a smaller scale.
- (3) AC, BD, BC, AD and CD have the same scale and AB has a smaller scale.
- (4) AC, BD, BC, AB and CD have the same scale and AD has a larger scale.
- (5) All six edges have the same scale.

To handle cases (1)–(3) we track two additional parameters, and show that the parity of one or both parameters is different for T and T' . In case (4) we again prove there is a change of parity, but of a third, more complicated parameter. For case (5) we use induction.

Figure 3a should make the definitions of the current paragraph clear. Orient the edges of the triangulation T so that the head of each edge has larger label. The *root edge* of T is the edge $\rho = (0, n-1)$. Now construct the following oriented tree \hat{T} . First, augment T by adding a vertex v to the unbounded face, and join it to all vertices of the polygon. Let \hat{T}_0 be the planar dual of the augmented graph; then \hat{T} is formed by removing all edges of \hat{T}_0 lying entirely within the unbounded face of T . For each edge e of T there is a unique edge \hat{e} of \hat{T} crossing e . Orient \hat{e} from the left to the right of e (when following e from tail to head). Root \hat{T} at the edge $\hat{\rho}$, whose head is the unique node of \hat{T} with out-degree zero. Note that \hat{T} is a tree, which we call the *dual tree* of T .

Given an edge $e = UV$ of T with $e \neq \rho$, the triangle containing the head of \hat{e} is incident to both U and V ; let W be its third node. Necessarily either $W < \min(V, U)$ or $W > \max(V, U)$. In the first case we say \hat{e} is a *left turn*, in the second we say it is a *right turn*.

Given a subgraph S of triangulation T , we let \hat{S} be the “dual” subgraph of \hat{T} , with the same vertex set as \hat{T} and with edge set

$$E(\hat{S}) = \{\hat{e} : e \in E(S)\};$$




Figure 3. The dual trees of an 8-gon and of a sub-triangulation of a 12-gon

this is illustrated in Figure 3b. A node of \hat{S} is a *leaf* if it has degree one. For each node v of \hat{S} , let $g_S(v)$ and $d_S(v)$ be defined as follows. Write r for the root (the unique node of out-degree 0) of the tree component of \hat{S} containing v . Then $g_S(v)$ and $d_S(v)$ are the number of left- and right-turns on the path from v to r , respectively; see Figure 4a.




Figure 4. In both subfigures, left-turn edges are red and right-turn edges are blue.

Recall that T' is obtained from T by flipping edge AC within quadrilateral $ABCD$. For each $1 \leq i \leq \lceil \log_r n \rceil$, let S_i be the subgraph of T with edge set $E(S_i) = \{e \in E(T) : \sigma_e = i\}$, and

let S'_i be the subgraph of T' with edge set $E(S'_i) = \{e \in E(T') : \sigma_e = i\}$. Then let \hat{S}_i and \hat{S}'_i be as defined in the preceding paragraph (so \hat{S}_i is a subgraph of \hat{T} and \hat{S}'_i is a subgraph of \hat{T}'). Let

$$G(T) = \sum_{i=1}^{\lceil \log_r n \rceil} \sum_{v \in V(\hat{S}_i)} g_{S_i}(v), \quad \text{and let} \quad D(T) = \sum_{i=1}^{\lceil \log_r n \rceil} \sum_{v \in V(\hat{S}_i)} d_{S_i}(v).$$

The following proposition implies that in cases (1), (2) and (3), flipping edge AC yields a change in parity of at least one of G and D .

Proposition 5. *If the scales of the edges AB , AC , AD , BC , BD and CD are as in cases (1), (2) or (3) above, then either $G(T') = G(T) - 1$ or $D(T') = D(T) + 1$ or both.*

Proof. Let $\sigma = \sigma_{AC}$. We first claim that for all $i \neq \sigma$, the contributions to $G(T)$ and to $D(T)$ from scale- i nodes are unchanged by the edge flip operation, i.e.,

$$\sum_{i \neq \sigma} \sum_{v \in V(\hat{S}_i)} g_{S_i}(v) = \sum_{i \neq \sigma} \sum_{v \in V(\hat{S}'_i)} g_{S'_i}(v) \quad \text{and} \quad \sum_{i \neq \sigma} \sum_{v \in V(\hat{S}_i)} d_{S_i}(v) = \sum_{i \neq \sigma} \sum_{v \in V(\hat{S}'_i)} d_{S'_i}(v). \quad (2)$$

We prove these equalities in case (1); the other two cases are similar but easier.

The triangle containing the head of the edge \hat{e}_{AB} dual to AB is ABC in T and is ABD in T' . The case (1) assumptions on the scales of the edges then imply that \hat{e}_{AB} has out-degree 0 and in-degree 1 in $\hat{S}_{\sigma_{AB}}$. In particular, it is the root of its component of $\hat{S}_{\sigma_{AB}}$. Moreover, \hat{e}_{AB} is a right-turn edge since C and D are both larger than A and B .

Similarly, the triangle containing the tail of the edge \hat{e}_{CD} dual to CD is ACD in T and is BCD in T' . The assumptions on the scales of edges again imply that \hat{e}_{CD} has in-degree 1 and out-degree 0 within $\hat{S}_{\sigma_{CD}}$, so is the root of its component of $\hat{S}_{\sigma_{CD}}$. Moreover, \hat{e}_{AB} is a left-turn edge since A and B are both smaller than C and D .

Since the structures of T and of T' are unaffected outside of the quadrilateral $ABCD$, the equalities in (2) follow in case (1).

We now restrict our attention to scale σ . We write $g(\cdot) = g_{S_\sigma}(\cdot)$ and $d(\cdot) = d_{S_\sigma}(\cdot)$, and likewise write $g'(\cdot) = g_{S'_\sigma}(\cdot)$ and $d'(\cdot) = d_{S'_\sigma}(\cdot)$. Note that all nodes not lying within the quadrilateral $ABCD$ either belong to both S_σ and S'_σ or belong to neither of S_σ and S'_σ .

The remainder of the proof boils down to inspection of Figures 5, 6 and 7. In case (1), observe (see Figure 5) that $g(u) = g'(u) = a + 1$ and $d(u) = d'(u) = b + 1$, which implies that $(g(q), g(q)) = (g'(q), d'(q))$ for all nodes q not lying within $ABCD$. Since $g(v) + g(x) = 2a + 1 = g(p) + g(z) + 1$ and $d(v) + d(z) = 2b = d(p) + d(z) - 1$, it follows that $G(T) = G(T') + 1$ and $D(T') = D(T) - 1$.

Figure 6 depicts the situation in case (2). In this case $d(u) = d'(u)$ and $d(y) = d'(y)$, which implies that $d(q) = d'(q)$ for all q not lying in $ABCD$. Since $d'(z) + d'(p) = d(v) + d(x) + 1$, it follows that $D(T') = D(T) + 1$.

Finally, Figure 7 relates to case (3). In this case $g(u) = g'(u)$, $g(y) = g'(y)$, and $g(v) + g(x) = g'(z) + g(p) + 1$, so the same logic as above implies that $G(T') = G(T) - 1$. This completes the proof. \square

We now turn our attention to cases (4) and (5). Consider any subgraph S of T , and let \hat{H} be a connected component of \hat{S} . Note that \hat{H} is a rooted sub-binary tree (i.e. every node has degree at most three; see Figure 3b). Let \tilde{H} be the tree obtained from \hat{H} as follows (see Figure 4a and 4b). First, if the root r of \hat{H} has exactly two children then add a new node \tilde{r} incident only to r and reroot at \tilde{r} . Next, suppress all nodes of degree exactly two to obtain a rooted binary tree. We call \tilde{H} the *reduced tree* of \hat{H} .

Proposition 6. *For all $1 \leq i \leq \lceil \log_r n \rceil$ and all components \hat{H} of \hat{S}_i , the reduced tree \tilde{H} has at most $2r - 1$ leaves.*

Proof. Fix any node u of \tilde{H} with in-degree zero, and consider the edge uv incident to u in \tilde{H} . Then uv is dual to an edge AB with $\sigma_{AB} = i$ so with $r^{i-1} < B - A \leq r^i$. Now fix another node w




Figure 5. The left-turn and right-turn labels near quadrilateral $ABCD$ in T and T' : case (1). Here $(g(x), d(x)) = (a, b)$, $(g(v), d(v)) = (a + 1, b)$ and $(g(u), d(u)) = (a + 1, b + 1)$.




Figure 6. The left-turn and right-turn labels near $ABCD$ in T and T' : case (2).




Figure 7. The left-turn and right-turn labels near $ABCD$ in T and T' : case (3).

of \tilde{H} with in-degree zero, write wx for the edge incident to w in \tilde{H} , and let CD be its dual edge. Then necessarily either $A < B < C < D$ or $C < D < B < A$.

Next consider an edge yr from a child of r to r in H . Writing EF for the edge dual to yr , then the observation of the preceding paragraph implies that $F - E > r^{i-1} \cdot \ell$, where ℓ is the number of nodes with in-degree zero in the subtree rooted at y . On the other hand, $\sigma_{EF} = i$ so $F - E \leq r^i$; so $\ell \leq r - 1$. If r has only one child (so is a leaf itself) this yields that \tilde{H} has at most r leaves. If r has two children then each of their subtrees contains at most $r - 1$ leaves; in this case \tilde{r} is also a leaf, so the total number of leaves is at most $2(r - 1) + 1$. \square

For any subgraph S of the triangulation T , the embedding of \hat{T} in the plane induces a total order of the connected components of \hat{S} , given by the order their roots are visited by a clockwise tour around the contour of \hat{T} starting from the head of the root edge ρ . For each $1 \leq i \leq \lceil \log_r n \rceil$, list the components of \hat{S}_i in the order just described as $H_{i,1}, \dots, H_{i,\ell}$, where $\ell = \ell(T, i)$ is the number of such components. Then, for $1 \leq j \leq \ell(T, i)$ let $\tilde{H}_{i,j}$ be the reduced tree of $H_{i,j}$.

Each tree from $(\tilde{H}_{i,j}, i \leq \lceil \log_r n \rceil, j \leq \ell(T, i))$ is a dual to a unique triangulation $\tilde{T}_{i,j}$ of a polygon, as in Figure 4b. Proposition 6 implies that $\tilde{T}_{i,j}$ belongs to an associahedron \mathcal{A}_k for some $k \leq 2r - 1$. Let ϕ be a proper colouring of the disjoint union of $(\mathcal{A}_k, k \leq 2r - 1)$, with colours $\{1, \dots, \chi(\mathcal{A}_{2r-1})\}$, and define

$$I(T) = \left(\sum_{i=0}^{\lceil \log_r n \rceil} \sum_{j=1}^{\ell(T,i)} \phi(\tilde{T}_{i,j}) \right) \pmod{\chi(\mathcal{A}_{2r-1})}.$$

Proposition 7. *In cases (4) and (5), we have $I(T) \neq I(T')$.*

Proof. We write $vx = \hat{e}$ for the dual edge of AC in T , and $zp = \hat{e}_{BD}$ for the dual edge of BD in T' . Figures 8 and 9 illustrate cases (4) and (5) respectively.

The clockwise contour exploration of a rooted plane tree is a walk around the outside of the tree which starts and finishes at the root, keeping the unbounded face to its left at all times. This walk traverses each edge exactly twice, and records the vertices it visits in sequence, with repetition. In cases (4) and (5), for the clockwise contour explorations of \hat{T} and of \hat{T}' , there are (possibly empty) strings P_1, \dots, P_5 so that the sequences recorded by the contour explorations of \hat{T} and of \hat{T}' are of the form

$$P_1 s x v w P_2 w v u P_3 u v x y P_4 y x s P_5$$

and

$$P_1 s p w P_2 w p z u P_3 u z y P_4 y z p s P_5,$$

respectively; again, see Figures 8 and 9.

The contour explorations of \hat{T} and \hat{T}' agree until they visit dual vertices lying within $ABCD$. It follows that if $H_{\sigma,j}$ is the component of S_σ containing \hat{e}_{AC} , then the component of S'_σ containing \hat{e}_{BD} is $H'_{\sigma,j}$.

In case (4), since x has two children in $H_{\sigma,j}$, by construction it is the unique child of the root of $\tilde{H}_{\sigma,j}$. It is thus natural to identify s with the root of $\tilde{H}_{\sigma,j}$. We may likewise identify s with the root of $\tilde{H}'_{\sigma,j}$, since p has two children in $H'_{\sigma,j}$. After the addition of s as a root, the nodes v, x, p and z all have degree 3, so none of these nodes are suppressed when constructing $\tilde{H}_{\sigma,j}$ and $\tilde{H}'_{\sigma,j}$ from $H_{\sigma,j}$ and $H'_{\sigma,j}$.

In case (5), nodes v, x, p and z all have degree 3, and the edges xs and ps belong to $H_{\sigma,j}$ and $H'_{\sigma,j}$, respectively.

It follows from the two preceding paragraphs that in cases (4) and (5), we may view v and x as nodes of both $H_{\sigma,j}$ and $\tilde{H}_{\sigma,j}$, and p and z as nodes of both $H'_{\sigma,j}$ and $\tilde{H}'_{\sigma,j}$. It is then clear that flipping the edge AC in the triangulation T corresponds precisely to flipping the corresponding edge in $\tilde{T}_{\sigma,j}$ to form $\tilde{T}'_{\sigma,j}$.

Since $\tilde{T}_{\sigma,j}$ and $\tilde{T}'_{\sigma,j}$ are related by a single edge flip, and ϕ is a proper colouring, it follows that $\phi(\tilde{T}_{\sigma,j}) \neq \phi(\tilde{T}'_{\sigma,j})$. Since all other components of S_σ , and more generally of each $(S_i, 1 \leq i \leq \lceil \log_r n \rceil)$, are unchanged when moving from T to T' , the result follows. \square




Figure 8. The structure near the quadrilateral $ABCD$ in T and T' case (4). The dashed edges have scale $\sigma_{AD} > \sigma$, all other edges of the triangulations shown in the figure have scale σ .




Figure 9. The structure near the quadrilateral $ABCD$ in T and T' in case (5). All edges of the triangulations shown in the figure have scale σ .

Proof of Theorem 1. Consider the graph G with vertex set \mathbb{N}^4 where (l, m, r, z) and (l', m', r', z') are adjacent if $|l' - l| + |m' - m| + |r' - r| + |z' - z| \leq 3$. This graph is 128-regular, so is 129-colourable. (We are not optimizing constants.) Let $\kappa : \mathbb{N}^4 \rightarrow \{1, \dots, 129\}$ be a proper colouring of G .

In all situations covered by Proposition 2, the vectors $(l_{T'}, m_{T'}, r_{T'}, z_{T'})$ and (l_T, m_T, r_T, z_T) are adjacent in G , so we have $\kappa((l_{T'}, m_{T'}, r_{T'}, z_{T'})) \neq \kappa((l_T, m_T, r_T, z_T))$.

In the situations covered by Proposition 4, we have $c(T) \neq c(T')$.

Let $g(T) = G(T) \pmod 2$ and $d(T) = D(T) \pmod 2$. By Proposition 5, in cases (1)–(3), either $g(T) \neq g(T')$ or $d(T) \neq d(T')$ or both. Finally, in cases (4) and (5), by Proposition 7 we have $I(T) \neq I(T')$.

It follows that the data $\kappa((l_T, m_T, r_T, z_T))$, $c(T)$, $g(T)$, $d(T)$, and $I(T)$ are sufficient to distinguish T from all its neighbours in \mathcal{A}_n . Therefore,

$$\chi(\mathcal{A}_n) \leq 129 \cdot \lceil 3 \log_r n \rceil \cdot 2 \cdot 2 \cdot \chi(\mathcal{A}_{2r-1}),$$

for any $r \geq 3$. Taking $r = 3$ yields $\chi(\mathcal{A}_{2r-1}) = \chi(\mathcal{A}_5) = 3$, since \mathcal{A}_5 is a 5-cycle. It follows that

$$\chi(\mathcal{A}_n) \leq 12 \cdot 129 \cdot \lceil 3 \log_3 n \rceil = O(\log n). \quad \square$$

Acknowledgements

This research was undertaken at the 2018 Bellairs Workshop on Graph Theory, held at the Bellairs Research Institute of McGill University.

References

- [1] Louis J. Billera, Paul Filliman, and Bernd Sturmfels. Constructions and complexity of secondary polytopes. *Adv. Math.*, 83(2):155–179, 1990. doi: 10.1016/0001-8708(90)90077-Z. 1
- [2] Cesar Ceballos and Vincent Pilaud. The diameter of type D associahedra and the non-leaving-face property. *European J. Combin.*, 51:109–124, 2016. doi: 10.1016/j.ejc.2015.04.006. 2
- [3] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. *Canad. Math. Bull.*, 45(4):537–566, 2002. doi: 10.4153/CMB-2002-054-1. 1
- [4] Ruy Fabila-Monroy, David Flores-Peñaloza, Clemens Huemer, Ferran Hurtado, Jorge Urrutia, and David R. Wood. On the chromatic number of some flip graphs. *Discrete Math. Theor. Comput. Sci.*, 11(2):47–56, 2009. URL <https://dmtcs.episciences.org/460>. 2
- [5] Christophe Hohlweg, Carsten E. M. C. Lange, and Hugh Thomas. Permutahedra and generalized associahedra. *Adv. Math.*, 226(1):608–640, 2011. doi: 10.1016/j.aim.2010.07.005. 1
- [6] Ferran Hurtado and Marc Noy. Graph of triangulations of a convex polygon and tree of triangulations. *Comput. Geom. Theory Appl.*, 13(3):179–188, 1999. 1
- [7] Anders Johansson. *Some results on colourings of graphs*. PhD thesis, University of Umeå, 1994. 2
- [8] Jean-Louis Loday and María O. Ronco. Hopf algebra of the planar binary trees. *Adv. Math.*, 139(2):293–309, 1998. doi: 10.1006/aima.1998.1759. 1
- [9] Vincent Pilaud and Francisco Santos. The brick polytope of a sorting network. *European J. Combin.*, 33(4):632–662, 2012. doi: 10.1016/j.ejc.2011.12.003. 1
- [10] Vincent Pilaud and Christian Stump. Brick polytopes of spherical subword complexes and generalized associahedra. *Adv. Math.*, 276:1–61, 2015. doi: 10.1016/j.aim.2015.02.012. 1
- [11] Lionel Pournin. The diameter of associahedra. *Adv. Math.*, 259:13–42, 2014. doi: 10.1016/j.aim.2014.02.035. 2
- [12] Daniel D. Sleator, Robert E. Tarjan, and William P. Thurston. Rotation distance, triangulations, and hyperbolic geometry. *J. Amer. Math. Soc.*, 1(3):647–681, 1988. doi: 10.2307/1990951. 2
- [13] James Dillon Stasheff. Homotopy associativity of H -spaces. I, II. *Trans. Amer. Math. Soc.*, 108:293–312, 1963. doi: 10.1090/s0002-9947-1963-0158400-5. 1
- [14] Dov Tamari. *Monoïdes préordonnés et chaînes de Malcev*. Thèse, Université de Paris, 1951. 1

(L. Addario-Berry) Department of Mathematics and Statistics, McGill University, Montréal, Québec, Canada
Email address: louigi.addario@mcgill.ca

(B. Reed) School of Computer Science, McGill University, Montréal, Canada. CNRS, Projet COATI, I3S (CNRS and UNS) UMR7271 and INRIA, Sophia Antipolis, France. Instituto Nacional de Matemática Pura e Aplicada (IMPA), Brasil. Visiting Research Professor, ERATO Kawarabayashi Large Graph Project, National Institute of Informatics, Japan.

Email address: breed@cs.mcgill.ca

(A. Scott) Mathematical Institute, University of Oxford, Oxford OX2 6GG, United Kingdom
Email address: scott@maths.ox.ac.uk

(D. R. Wood) School of Mathematical Sciences, Monash University, Melbourne, Australia
Email address: david.wood@monash.edu