

GROWING RANDOM 3-CONNECTED MAPS

OR COMMENT S'ENFUIR DE L'HEXAGONE

L. ADDARIO-BERRY

Abstract. We use a growth procedure for binary trees [10], a bijection between binary trees and irreducible quadrangulations of the hexagon [6], and the classical angular mapping between quadrangulations and maps, to define a growth procedure for maps. The growth procedure is local, in that every map is obtained from its predecessor by an operation that only modifies vertices lying on a common face with some fixed vertex. As $n \rightarrow \infty$, the probability that the n 'th map in the sequence is 3-connected tends to $2^8/3^6$. The sequence of maps has an almost sure limit G_∞ , and we show that G_∞ is the *distributional* local limit of large, uniformly random 3-connected graphs.

1. Introduction

Here is a common situation in probability. We have found a result of the form "as $n \rightarrow \infty$, $X_n \rightarrow X_\infty$ in distribution", where $(X_n, 1 \leq n \leq \infty)$ are some sort of random objects. The Skorohod representation theorem then guarantees (in great generality) the existence of a coupling such that $X_n \rightarrow X_\infty$ almost surely. This result, though useful, is existential, and often the discovery of an *explicit* coupling leads to a deeper understanding of both the limit object and its finite approximations.

Given the substantial recent interest in random planar maps having the Brownian map as their (known or conjectural) scaling limit, it seems natural to seek such a coupling for random maps. (We hereafter refer to such a coupling as a *growth procedure*.) To date, however, all convergence results for such random planar maps have been distributional in nature. The goal of this note is to provide an explicit, local -- in the sense described in the abstract -- growth procedure for random planar maps. The result is a sequence of rooted maps $(M_n, 1 \leq n \leq \infty)$, such that $M_n \xrightarrow{\text{a.s.}} M_\infty$ as $n \rightarrow \infty$, by which we mean that balls of any fixed radius around the root almost surely stabilize.

We briefly summarize the structure and the arguments of the paper, then head right to details. We begin by considering *irreducible quadrangulations of the hexagon*: these are rooted maps with a single face of degree 6 and all other faces of degree 4, such that every cycle of length 4 bounds a face. (The root is a uniformly random oriented edge; it need not lie along the hexagonal face.) Such maps are in bijective correspondence with rooted binary plane trees [6]; we describe the bijection in Section 3.1. In Section 3.2 we describe how growing a binary tree -- transforming a degree-one vertex into a degree-three vertex -- changes the corresponding irreducible quadrangulation of the hexagon.

A growth procedure is already known to exist for random binary plane trees [10]; the almost sure limit T_∞ is a critical binary Galton-Watson tree, conditioned to survive. In Section 4 we use the bijection and properties of T_∞ to show that the corresponding growth procedure for irreducible quadrangulations of the hexagon has an almost sure limit. We then show, in Section 5, that the structure of the irreducible quadrangulations near their root asymptotically decouples from the structure near the hexagonal face. We use this fact together with the angular mapping between

Date: February 10, 2014.

2010 Mathematics Subject Classification. 60C05,60J80,05C10.

Key words and phrases. Random maps, random trees, random planar graphs, growth procedures.

quadrangulations and general maps to define a growth procedure $(G_n, 1 \leq n \leq \infty)$ for random maps. We show that the almost sure limit G_∞ is almost surely 3-connected and is the distributional limit of large random 3-connected maps. We conclude in Section 6 with a number of remarks and suggestions for future research.

2. Definitions

We briefly review some basic concepts regarding (planar) maps; more details can be found in, e.g., [8]. All our maps are connected. Also, maps are by default embedded in \mathbb{R}^2 rather than on the sphere \mathbb{S}^2 . For any graph or map M , we write $v(M)$ and $e(M)$ for the nodes and edges of M , respectively. The *corners* incident to a face f of M are the angles $\kappa = (\{u, v\}, \{v, w\})$ formed by consecutive edges along the face. We say κ is incident to f , to v , and to its constituent edges, and write $v(\kappa) = v$. The degree of f is the number of corners incident to f . Write $\mathcal{C}(M)$ for the set of corners of M .

For $\{u, v\} \in e(M)$, we use both (u, v) and uv as notation for the orientation of $\{u, v\}$ with tail u and head v . An oriented circuit C in M is *clockwise* if the bounded region of $\mathbb{R}^2 \setminus C$ lies to the right of C , and is otherwise counterclockwise. A *rooted map* is a pair $M = (M, vw)$, where $\{v, w\} \in e(M)$; vw is the *root edge*. The *root corner* $\rho = \rho(M)$ is the unique corner incident to v whose second incident edge is $\{v, w\}$, and the *root node* is $r(M) = u$.

Given a rooted map $M = (M, uv)$ and $R > 0$, write $M^{<R}$ for the submap induced by the set of nodes at graph distance less than R from u . We say a sequence $(M_n, n \geq 1)$ of (finite or infinite) locally finite rooted maps converges to rooted map M_∞ , and write $M_n \rightarrow M_\infty$, if for all $R > 0$, there is $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, $M_n^{<R}$ and $M_\infty^{<R}$ are isomorphic as rooted maps. Note that since the maps M_n are locally finite, so is their limit M_∞ . This is often called *local weak convergence*; see [2, 3, 7] for more details.

A *quadrangulation* is a map in which every face has degree four. In a *quadrangulation of the hexagon*, the unbounded face has degree six and all others have degree four. A quadrangulation, or a quadrangulation of the hexagon, is called *irreducible* if every cycle of length four bounds a face. It is easily seen that an irreducible quadrangulation of the hexagon is necessarily simple, and also 3-connected, so has a unique embedding by Whitney's theorem.

The definitions of the coming paragraphs are illustrated in Figure 1. A *plane tree* is a rooted tree $T = (T, r(T))$ together with an ordering of the children of each node v as $c_1(v), \dots, c_{k_T(v)}$, where $k_T(v)$ is the number of children of v . This collection of orderings uniquely specifies T as a planar map, with root $e^\rightarrow = e^\rightarrow(T) = (r(T), c_1(r(T)))$. Conversely, the orderings may be recovered from the embedding and the root edge $e^\rightarrow(T)$.

Viewed as a map, T has a unique (unbounded) face. Writing $n = |v(T)|$, list the (oriented) edges of this face in the order they are traversed by a counterclockwise tour¹ of T starting from $e^\rightarrow(T)$, as $e_1^\rightarrow(T), \dots, e_{2n-2}^\rightarrow(T)$, or simply $e_1^\rightarrow, \dots, e_{2n-2}^\rightarrow$ when the tree is clear from context. Write e_1, \dots, e_{2n-2} for the corresponding unoriented edges. Note that each edge of T appears exactly twice in this sequence. It is also convenient to set $e_{2n-1}^\rightarrow = e_1^\rightarrow$. We then have $\rho(T) = \{e_{2n-2}, e_1\}$.





Figure 1. A binary (plane) tree (T, r) . With labels as in the figure, $\kappa_3(v) \prec \rho \prec \kappa_1(v)$.

¹Recall that such a tour keeps the unbounded face to its left.

We now form a map from T as follows. By *closing* a bud corner $\kappa \notin \Delta$, we mean identifying $v(\kappa)$ and $v(\sigma(\kappa))$ to form an oriented edge $e_\kappa = (\alpha(v(\kappa)), v(\sigma(\kappa)))$, in such a way that the oriented cycle C_κ formed by following e_κ , then returning to $\alpha(v(\kappa))$ via $\llbracket v(\sigma(\kappa)), \alpha(v(\kappa)) \rrbracket$, is clockwise (the bounded region of $\mathbb{R}^2 \setminus C$ lies to its right).

Draw a hexagon so that T lies in its bounded face (interior), and label its interior corners $0, 1, \dots, 5$ so that for $\kappa \in \Delta$, $\sigma(\kappa)$ is a corner of the hexagon. (Later, we will also view the nodes of the hexagon as having labels $0, 1, 2, 3, 4, 5$, in the obvious way.) For each bud corner $\kappa \notin \Delta$, close κ (see Figure 3a). For each $\kappa \in \Delta$, identify $v(\kappa)$ with the hexagon corner labelled $\sigma(\kappa)$ (see Figure 3b).

Writing e^\rightarrow for the image of the root edge $e^\rightarrow(T)$ in M , the resulting map $M = (M, e^\rightarrow)$ is an irreducible, edge-rooted quadrangulation of a hexagon. The key point of this section (Proposition 1, below) is that the function taking T to M is bijective [6]. We use this fact more-or-less as a black box; however, the next two paragraphs contain a very brief sketch of one aspect of its proof, for the interested reader. A more detailed, very readable explanation can be found in [6, Section 4].



(a) Blue edges denote closures to internal corners.

(b) Red edges denote closures of corners in Δ .

(c) The induced orientation (edges without arrows are doubly oriented).

The map M inherits the orientations $\{e_\kappa, \kappa \in \mathcal{B}_B(T)\}$, which orients a subset of its edges (the orientation of e^\rightarrow need not respect the inherited orientation). The remaining edges of M are the edges of the hexagon, together with those edges of T joining internal nodes of T , and we view all these edges as doubly oriented, or oriented in both directions (see Figure 3c; in that figure the edge e^\rightarrow is indicated by a solid black arrow, whereas the inherited orientation is shown with empty white arrows). We say a doubly oriented edge is an outgoing edge from both its endpoints. Since T is binary, it follows that each non-hexagon node v of M has exactly three outgoing edges; such an orientation is called a *tri-orientation* of M . More precisely, a tri-orientation of M is an orientation of the edges of M (with both singly and doubly oriented edges permitted) such that every non-hexagon node has exactly three outgoing edges, and hexagon nodes have exactly two outgoing edges.

Finally, the "clockwise" orientation of the closure operation straightforwardly implies that M has no counterclockwise cycle in its interior, in the sense that any oriented cycle C in M containing at least one non-hexagon node and with its unbounded face to its right, must traverse some (singly) oriented edge of M from head to tail. It turns out that for any irreducible quadrangulation of the hexagon there is an *unique* tri-orientation of M with no counterclockwise cycle in its interior [6, Theorem 4.4]. Furthermore, the doubly-oriented edges of this tri-orientation form a hexagon plus a spanning tree T of the internal vertices of M , and T is a binary tree whose closure is M . Likewise, applying the closure operation to any tree T yields a map M whose "opening" is again T . It follows that the closure operation is a bijection.

Let \mathcal{T}_n be the set of binary trees with n internal nodes. Also, let \mathcal{M} be the set of pairs (M, e^\rightarrow) , where M is an irreducible quadrangulation of the hexagon and e^\rightarrow is an oriented edge of M at least one of whose endpoints is a non-hexagon node, and let $\mathcal{M}_n = \{(M, e^\rightarrow) \in \mathcal{M} : |v(M)| = n+6\}$.

Proposition 1 ([6], Theorems 4.7 and 4.8). *For each $n \geq 1$, the closure operation is a bijection between \mathcal{T}_n and \mathcal{M}_n .*

In the next subsection, we explain the effect of “growing” T -- transforming a bud into an internal node -- on the map M resulting from the closure operation.

3.2. Bud growth and map growth. Given a binary tree $T = (T, r(T))$ and a bud corner κ of T , *growing* T at κ means adjoining two buds incident to $v = v(\kappa)$, so κ becomes an internal node of degree three. Write T^+ for the resulting tree, which is still binary. Let M and M^+ be the closures of T and T^+ , respectively. Figures 4a and 4b depict the corresponding difference between M and M^+ ; this difference is *local*, in that it is confined to faces incident to $v(\sigma_T(\kappa))$. We now explain the transformation in detail.




Figure 4. Gray nodes are nodes of T but not of M (left), or of T^+ but not of M^+ (right). Blue edges have a thick part and a thin part, divided by a gray node; the thick part of each blue edge is itself an edge of T (left) or T^+ (right).

Let κ' be the first bud corner with $\sigma_T(\kappa') = \sigma_T(\kappa)$, in the sense that if κ^* is any other bud corner with $\sigma_T(\kappa^*) = \sigma_T(\kappa)$ then $\kappa' \prec \kappa^* \prec \kappa$. Then let $K(\kappa) = \{\kappa^* \in \mathcal{C}_B(T) : \sigma_T(\kappa') = \sigma_T(\kappa)\}$. In Figure 4a, the nodes incident to corners in $K(\kappa)$ are precisely the greyed nodes

Write $x = v(\sigma_T(\kappa))$, let $e_\ell = \{\alpha(v(\kappa')), x\}$ and let $e_r = \{\alpha(v(\kappa)), x\}$, and let f_ℓ and f_r be the faces of M lying to the left and right of e_ℓ and e_r , respectively. Then let v_ℓ and v_r be the vertices of M diagonally opposite $\alpha(v(\kappa'))$ and $\alpha(v(\kappa))$ on f_ℓ and on f_r , respectively, and let κ_ℓ and κ_r be the corners of f_ℓ and f_r incident to v_ℓ and v_r . Again, see Figure 4a.

With these definitions, M^+ is formed from M as follows. For all $\xi \in K(\kappa)$ remove e_ξ from M . This creates a face of degree $2|K(\kappa)| + 4$. Add a vertex v in this face (recall that we also wrote $v = v(\kappa)$; this is deliberate), and add edges from v to $\alpha(v(\xi))$ for each $\xi \in K(\kappa)$, and from v to $v(\kappa_\ell)$ and $v(\kappa_r)$.

To prove that this description is valid, argue as follows.² Viewing T as a subtree of T^+ in the natural way, for $\kappa' \in \mathcal{C}(T) \setminus \{\kappa\}$ we have $S_T(\kappa') = S_{T^+}(\kappa')$. It follows that if $\kappa' \in \mathcal{C}(T) \setminus K(\kappa)$ then $\sigma_T(\kappa') = \sigma_{T^+}(\kappa')$. In T^+ the corner $\kappa_1(v)$ is an internal corner, and has $S_{T^+}(\kappa_1(v)) = S_T(\kappa)$. Furthermore, for all $\kappa' \in K(\kappa) \setminus \{\kappa\}$, in T^+ we have $\kappa' \prec \kappa_1(v) \prec \sigma_T(\kappa') = \sigma_T(\kappa)$ and thus $\sigma_{T^+}(\kappa') = \kappa_1(v)$.

Next write ξ_ℓ and ξ_r for the bud corners incident to the left and right children of v in T^+ (these are the two greyed nodes incident to v in Figure 4b). Then $S_{T^+}(\xi_\ell) = S_T(\kappa) - 1$ and $S_{T^+}(\xi_r) =$

²If Figure 4 is sufficiently convincing, feel free to skip straight to Section 4.

$S_{\mathbb{T}}(\kappa) + 1$. We claim that $\sigma_{\mathbb{T}^+}(\xi_\ell) = \kappa_\ell$ and $\sigma_{\mathbb{T}^+}(\xi_r) = \kappa_r$; proving this will establish the validity of the above description.

In proving the above claim, it is useful to extend the domain of definition of $\sigma_{\mathbb{T}}$ from $\mathcal{C}_B(\mathbb{T})$ to $\mathcal{C}(\mathbb{T})$ as follows. For $\kappa \in \mathcal{C}_I(\mathbb{T})$, if the first corner κ' after κ in the counterclockwise tour around \mathbb{T} is internal set $\sigma_{\mathbb{T}}(\kappa) = \kappa'$, and otherwise set $\sigma_{\mathbb{T}}(\kappa) = \sigma_{\mathbb{T}}(\kappa')$. We likewise define $\sigma_{\mathbb{T}^+}(\kappa)$ for $\kappa \in \mathcal{C}_I(\mathbb{T}^+)$. With the above definition, for all $\kappa \in \mathcal{C}_I(\mathbb{T})$, $S_{\mathbb{T}}(\sigma_{\mathbb{T}}(\kappa)) = S_{\mathbb{T}}(\kappa) - 6\mathbf{1}_{[\kappa \prec \rho \prec \sigma_{\mathbb{T}}(\kappa)]} - 1$, and all corners κ' with $\kappa \prec \kappa' \prec \sigma_{\mathbb{T}}(\kappa)$ have $S_{\mathbb{T}}(\kappa') \geq S_{\mathbb{T}}(\kappa) - 6\mathbf{1}_{[\kappa \prec \rho \preceq \kappa']}$. The analogous assertion holds for \mathbb{T}^+ .

First consider ξ_ℓ ; we must show that κ_ℓ is the first corner κ' after ξ_ℓ in \mathbb{T}^+ with $S_{\mathbb{T}^+}(\kappa') = S_{\mathbb{T}^+}(\xi_\ell) - 6\mathbf{1}_{[\xi_\ell \prec \rho \preceq \kappa']}$ $= S_{\mathbb{T}}(\kappa) - 1 - 6\mathbf{1}_{[\xi_\ell \prec \rho \preceq \kappa']}$. All corners $\kappa' \in \mathcal{C}(\mathbb{T}^+)$ with $\xi_\ell \prec \kappa' \preceq \sigma_{\mathbb{T}}(\kappa)$ have $S_{\mathbb{T}^+}(\kappa') \geq S_{\mathbb{T}}(\kappa) - 6\mathbf{1}_{[\kappa \prec \rho \preceq \kappa']}$; this follows from the definition of $S_{\mathbb{T}^+}$ for the corners of $\mathcal{C}(\mathbb{T}^+) \setminus \mathcal{C}(\mathbb{T})$, and for the remaining corners follows from the definition of $\sigma_{\mathbb{T}}(\kappa)$. Also, $\sigma_{\mathbb{T}}(\kappa)$ is internal and $\kappa_\ell = \sigma_{\mathbb{T}}(\sigma_{\mathbb{T}}(\kappa))$, so all corners κ' with $\sigma_{\mathbb{T}}(\kappa) \prec \kappa' \prec \kappa_\ell$ have $S_{\mathbb{T}^+}(\kappa') \geq S_{\mathbb{T}^+}(\sigma_{\mathbb{T}}(\kappa)) - 6\mathbf{1}_{[\sigma_{\mathbb{T}}(\kappa) \prec \rho \preceq \kappa']}$. Since ρ can not lie both between κ and $\sigma_{\mathbb{T}}(\kappa)$ and between $\sigma_{\mathbb{T}}(\kappa)$ and κ_ℓ , the result follows.

The argument for ξ_r is similar so we only sketch it. Let ξ^* be the corner of f_r incident to $\alpha(\kappa)$, and note that ξ^* is a corner of both \mathbb{T} and \mathbb{T}^+ . Thus $S_{\mathbb{T}^+}(\xi^*) = S_{\mathbb{T}}(\xi^*) = S_{\mathbb{T}}(\kappa) + 3$. Since $\kappa_r = \sigma_{\mathbb{T}}(\sigma_{\mathbb{T}}(\xi^*))$, by twice applying the definition of the attachment corner, κ_r must be the first corner κ' after ξ_r in \mathbb{T}^+ with $S_{\mathbb{T}^+}(\kappa') \leq S_{\mathbb{T}}(\kappa) + 1 - 6\mathbf{1}_{[\xi_r \prec \rho \preceq \kappa']}$ $= S_{\mathbb{T}^+}(\xi_r) - 6\mathbf{1}_{[\xi_r \prec \rho \preceq \kappa']}$.

4. Growing uniformly random trees and, thus, maps

In the infinite rooted binary tree \mathbb{T} every node has precisely two children -- one left and one right child -- so all nodes have degree three but the root, which has degree two. Depth- n nodes in \mathbb{T} may be represented as strings in $\{-1, 1\}^n$, with -1 and 1 representing left and right. We adopt this point of view, and identify \mathbb{T} with its node set $\bigcup_{i \geq 0} \{-1, 1\}^i$, where $\{-1, 1\}^0 = \{\emptyset\}$ and \emptyset is the root of \mathbb{T} .

It is temporarily useful to view binary trees as subtrees of \mathbb{T} as follows. For a given binary tree \mathbb{T} , add a vertex z in the middle of edge $e^{\rightarrow}(\mathbb{T})$. Identify z with $r(\mathbb{T})$, and the head and tail of e^{\rightarrow} with the left and right children of $r(\mathbb{T})$, respectively. Recursively embed the remaining nodes by making first and second children in \mathbb{T} respectively correspond to left and right children in \mathbb{T} . See Figure 5 for an illustration. The plane tree \mathbb{T} can be recovered from its representation as a subtree of \mathbb{T} -- essentially by replacing the path from 1 to 0 through \emptyset by a single edge -- so this viewpoint is reasonable. With this perspective we have $e^{\rightarrow}(\mathbb{T}) = (1, -1)$ and $r(\mathbb{T}) = 1$; as usual $e^{\rightarrow}(\mathbb{T})$ is the second edge incident to the root corner ρ .

The following result of Luczak and Winkler [10] is key tool in the current work.

Theorem 2 ([10], Theorem 4.1). *There exists a sequence $(\mathbb{T}_n, n \geq 1)$ of random binary trees with the following properties.*

- (1) For each $n \geq 1$, \mathbb{T}_n is uniformly distributed in \mathcal{T}_n .
- (2) For each $n \geq 1$, there is a bud corner ξ_n of \mathbb{T}_n such that \mathbb{T}_{n+1} is obtained from \mathbb{T}_n by growing at ξ_n . In particular, the sequence is increasing so has a limit $\mathbb{T}_\infty \subset \mathbb{T}$.
- (3) The limit \mathbb{T}_∞ is a critical binary Galton-Watson tree, conditioned to be infinite.

Here is a more detailed explanation of property (3). Let $(X_n, n \geq 1)$ be iid with $\mathbf{P}\{X_n = 1\} = 1/2 = \mathbf{P}\{X_n = -1\}$, and for each n let $Y_n = -X_n$. Let P be the infinite path $\{(X_1, \dots, X_i), i \geq 0\}$ in \mathbb{T} . Let $(B_n, n \geq 1)$ be independent Galton-Watson trees with offspring distribution μ ,

where $\mu(\{0\}) = 1/2 = \mu(\{2\})$, and for each $i \geq 1$, append B_i to P by rooting B_i at node $(X_1, \dots, X_{i-1}, Y_i)$. The resulting tree (see Figure 6 for an illustration) has the law of \mathbb{T}_∞ .




Figure 5. *Left*: a part of \mathbb{T} , with some node labels shown. *Center*: a binary tree \mathbb{T} . *Right*: \mathbb{T} viewed as a subtree of \mathbb{T} .

By Proposition 1, the closure operation associates to each tree T_n a map M_n which is uniformly distributed in \mathcal{M}_n . It therefore seems reasonable to expect that applying the closure rules to \mathbb{T}_∞ yields a map M_∞ and that $M_n \xrightarrow{\text{a.s.}} M_\infty$. This is indeed the case, and proving so is the subject of the remainder of the section.

We now view \mathbb{T}_∞ as a binary plane tree (rather than as a subtree of \mathbb{T}). The set of corners $\mathcal{C}(\mathbb{T}_\infty)$ consists pairs (e, e') , where e' follows e in the counterclockwise walk around (the unique, infinite face of) \mathbb{T}_∞ .³ Write \prec for the total order on $\mathcal{C}(\mathbb{T}_\infty)$ given by this walk.

The bud corners $\mathcal{C}_B(\mathbb{T}_\infty)$ and internal corners $\mathcal{C}_I(\mathbb{T}_\infty)$ are defined as before. Write \mathcal{C}^ℓ and \mathcal{C}^r for the set of corners following and preceding ρ in the counterclockwise walk around \mathbb{T}_∞ , respectively. Define labels $S_{\mathbb{T}_\infty} : \mathcal{C}(\mathbb{T}_\infty) \rightarrow \mathbb{Z}$ exactly as in (1). The second description of the labels, given just after (1) for finite trees, again applies: $S_{\mathbb{T}_\infty}(\rho) = -4\mathbf{1}_{[\rho \text{ is a bud corner}]}$, and in a counterclockwise walk, labels decrease by one when leaving an internal corner and increase by three when leaving a bud corner (except when the walk arrives at the root corner; then one must additionally subtract six).

For $\kappa \in \mathcal{C}_B(\mathbb{T}_\infty)$, let $\sigma(\kappa) = \sigma_{\mathbb{T}_\infty}(\kappa)$ be the first corner κ' following κ in the counterclockwise walk around \mathbb{T}_∞ for which $S_{\mathbb{T}_\infty}(\kappa') \leq S_{\mathbb{T}_\infty}(\kappa) - 6\mathbf{1}_{[\kappa \prec \rho \preceq \kappa']}$, if such a corner exists. Otherwise, set $\sigma_{\mathbb{T}_\infty}(\kappa) = -\infty$. It is immediate that if $\sigma_{\mathbb{T}_\infty}(\kappa) \neq -\infty$ then $S_{\mathbb{T}_\infty}(\sigma(\kappa)) = S_{\mathbb{T}_\infty}(\kappa) - 6\mathbf{1}_{[\kappa \prec \rho \preceq \sigma(\kappa)]}$. The set of corners κ with $\sigma(\kappa) = -\infty$ is the analogue of the set Δ of corners attaching to the hexagon when closing a finite binary tree. The next proposition states that this set vanishes in the $n \rightarrow \infty$ limit, and is my excuse for the paper's subtitle.

Proposition 3. *There are almost surely no corners $\kappa \in \mathcal{C}(\mathbb{T}_\infty)$ with $\sigma_{\mathbb{T}_\infty}(\kappa) = -\infty$.*

Proof. For any corner $\xi \in \mathcal{C}(\mathbb{T}_\infty)$, all but finitely many elements of \mathcal{C}^ℓ follow ξ in the walk. It thus suffices to show that $\inf\{S_{\mathbb{T}_\infty}(\kappa) : \kappa \in \mathbb{T}_\infty\} = -\infty$.

View \mathbb{T}_∞ as built from the path P , the random variables X_i and the random trees B_i as above, and for $n \geq 0$ let v_n be the n 'th node along P (so $v_0 = r(\mathbb{T}_\infty)$). Then for all $n \geq 1$, $\kappa_1(v_n) \in \mathcal{C}^\ell$ and $S_{\mathbb{T}_\infty}(\kappa_1(v_n)) = \sum_{i=1}^n X_i$. Thus $(S_{\mathbb{T}_\infty}(\kappa_1(v_n)), n \geq 1)$ forms a symmetric simple random walk and so $\inf\{S_{\mathbb{T}_\infty}(\kappa_1(v_n)), n \geq 1\} = -\infty$ almost surely. \square




Figure 6. A part of \mathbb{T}_∞ , with its unique infinite path in purple.

³We call it a walk rather than a tour since it is not closed.

For $\kappa \in \mathcal{C}_B(\mathbb{T}_\infty)$ define the closure operation, the closure edge e_κ , and the oriented cycle C_κ exactly as in Section 3.1. By the minimality of $\sigma(\kappa)$, all corners lying κ' within the bounded face of C_κ have $S_{\mathbb{T}_\infty}(\kappa') > S_{\mathbb{T}_\infty}(\kappa) - 6$. Since any infinite path leaving κ follows P for all but finitely many steps, and the corners along P take unboundedly large negative values, it follows that the interior of C_κ contains only finitely many vertices of \mathbb{T}_∞ .

Let M_∞ be formed by closing κ in the corner $\sigma(\kappa)$ for each $\kappa \in \mathcal{C}_B(\mathbb{T}_\infty)$.

Proposition 4. M_∞ is almost surely locally finite.

Proof. First observe that for $\kappa, \xi \in \mathcal{C}_B(\mathbb{T}_\infty)$, if $\kappa \prec \xi \prec \sigma(\kappa)$ then $\kappa \prec \sigma(\xi) \preceq \sigma(\kappa)$. It follows that if $\xi \prec \kappa$ then either $\sigma(\xi) \prec \kappa$ or $\sigma(\kappa) \prec \sigma(\xi)$. Since \mathbb{T}_∞ is a.s. locally finite and the cycles C_κ a.s. have finite interior, it suffices to show that a.s. for all internal corners ξ of \mathbb{T}_∞ , the set $\{\kappa \in \mathcal{C}_B(\mathbb{T}_\infty) : \sigma(\kappa) = \xi\}$ is a.s. finite. Let $\pi(\xi)$ be the maximal corner ξ' (with respect to \prec) preceding ξ for which $S_{\mathbb{T}_\infty}(\xi') < S_{\mathbb{T}_\infty}(\xi) - 6$.

All but finitely many corners preceding ξ lie in \mathcal{C}_r , and a similar argument to that above shows that the corners along P lying in \mathcal{C}_r take unboundedly large negative values. Since a.s. only finitely many corners lie between any two corners of $\mathcal{C}(\mathbb{T}_\infty)$ with respect to \prec , it follows that $\pi(\xi)$ is well-defined. Furthermore, $\pi(\xi)$ must be a bud corner since its \prec -successor has a larger label than its own. It follows that $\xi \prec \sigma(\pi(\xi))$. Thus, by the observation from the start of the first paragraph, if κ is any bud corner with $\sigma(\kappa) = \xi$ then necessarily $\pi(\xi) \prec \kappa \prec \xi$; and there are only finitely many such corners. \square

Corollary 5. $M_n \xrightarrow{\text{a.s.}} M_\infty$ as $n \rightarrow \infty$.

Proof. Recall that \mathbb{T}_{n+1} is obtained from \mathbb{T}_n by growing at corner ξ_n , and the description in Section 3.2 of how such growth transforms the associated map. It follows from this description that for $\kappa \in \mathcal{C}_B(\mathbb{T}_n)$, if $\sigma_{\mathbb{T}_{n+1}}(\kappa) \neq \sigma_{\mathbb{T}_n}(\kappa)$ then $\sigma_{\mathbb{T}_{n+1}}(\kappa) = (\{\alpha(v(\xi_n)), v(\xi_n)\}, \{v(\xi_n), c_1(v(\xi_n))\})$. Since $c_1(v(\xi_n))$ is not a node of \mathbb{T}_n , it follows that $\sigma_{\mathbb{T}_{n+1}}(\kappa) \notin \mathcal{C}_B(\mathbb{T}_n)$.

Now, for each corner $\kappa \in \mathcal{C}(\mathbb{T}_\infty)$, let $\tau(\kappa) = \inf\{m : \kappa \in \mathcal{C}(\mathbb{T}_m)\}$; then $\tau(\kappa)$ is almost surely finite. By the fact in the preceding paragraph, for all $n \geq \max(\tau(\kappa), \tau(\sigma_{\mathbb{T}_\infty}(\kappa)))$, we have $\sigma_{\mathbb{T}_n}(\kappa) = \sigma_{\mathbb{T}_\infty}(\kappa)$. Since M_∞ is a.s. locally finite, it follows that for any $R > 0$, there is an a.s. finite time n_0 such that for all $m, n \geq n_0$, $M_n^{<R}$ and $M_m^{<R}$ are isomorphic. \square

5. Three-connected maps

Given a locally finite map G , the *angular mapping* associates to G a quadrangulation Q as follows. Add a vertex v_f in each face f of G . For each corner κ incident to f , add an edge between v_f and $v(\kappa)$ attaching in corner κ . Then erase the edges of G (see Figure 7). Here consider G and Q as embedded in \mathbb{S}^2 (in \mathbb{R}^2 there is a choice of how to draw the edges of Q lying in the unbounded face of G), but in \mathbb{S}^2 this choice vanishes. Then the angular mapping is a 2-to-1 map; its inverse images may be found by properly 2-coloring the vertices of Q , and choosing one of the two colour classes to form the nodes of G . Note that the number of edges of G is the number of faces of Q .

There is a natural function φ taking oriented edges of G to oriented edges of Q : for uv an oriented edge of G , let f be the face of G lying to the right of G , and let $\varphi(uv) = uv_f$. Note that the tail of $\varphi(uv)$ is always a node of G . This yields an extension of the angular mapping to rooted maps, which sends (G, e) to $(Q, \varphi(e))$. The mapping is now bijective, since the orientation of $\varphi(e)$ determines which of the colour classes of Q forms the nodes of G .

By Proposition 1, for any irreducible quadrangulation of a hexagon $M \in \mathcal{M}$, we may view M as arising from a binary tree T by the closure bijection. We may thus canonically label the nodes

of the hexagonal face of M with labels $0, 1, 2, 3, 4, 5$. We hereafter view all $M \in \mathcal{M}$ as endowed with such labels, and for $i \in \{0, 1, 2, 3, 4, 5\}$ write $M(i)$ for the rooted quadrangulation obtained from M_n by adding an edge $e^* = \{i, i + 3 \pmod 6\}$ from i to the diagonally opposing node of the hexagon.

Now let $(M_n, 1 \leq n \leq \infty)$ be as in Section 4. Independently of $(M_n, 1 \leq n \leq \infty)$, let U be uniformly distributed in $\{0, 1, 2, 3, 4, 5\}$. For $1 \leq n \leq \infty$, let $Q_n = M_n(U)$. Then Q_n is a rooted quadrangulation, and we let G_n be the associated rooted map under the inverse of the angular mapping.

It is well-known (see, e.g., [6, Theorem 3.1]) that if a quadrangulation M is associated with a map G under the angular mapping, then G is 3-connected if and only if M is irreducible. The quadrangulation Q_n need not be irreducible, since there may be a 3-edge path between the endpoints of e^* passing through the interior of the hexagon in M_n . However, it turns out that Q_n is irreducible with uniformly positive probability, and hence G_n is 3-connected with uniformly positive probability. The main point of this section is to show (in Proposition 6) that G_n a.s. converges, and to identify the limit (in Theorem 7) as "the uniform infinite 3-connected planar map" (or -- thanks to Whitney's theorem -- graph). However, we first briefly describe how the growth dynamics modify G_n (though this information is not in fact needed in the paper).




Figure 7. A rooted map (with grey nodes and black edges) and its image under the angular mapping (with white and grey nodes and blue edges).

The effect on G_n of growing T_n depends on whether the growing bud corner is attached to a primal or a facial vertex. If it is attached to a facial vertex then growing T_n adds an edge within the face. If it is attached to a primal vertex then growing instead "uncontracts" the primal vertex, turning it into two vertices joined by an edge; this may be seen as adding an edge to the facial dual graph of G_n . In either case, this is a "local" modification in that it only affects the nodes that lie on a common face with a given vertex. We now turn to the main business of the section.

Proposition 6. $G_n \xrightarrow{\text{a.s.}} G_\infty$ as $n \rightarrow \infty$.

Proof. Fix any map G and let M be the image of G under the angular mapping. Then for all $\{u, v\} \in e(G)$, there exists a path of length two between u and v in Q . Thus, for all $x, y \in v(G)$, $d_G(x, y) \geq d_M(x, y)/2$, where d_G and d_Q denote graph distance in G and Q respectively. Since Q_n and G_n have the same root node and $Q_n \xrightarrow{\text{a.s.}} Q_\infty$, the result follows. \square

Theorem 7. Let \hat{G}_n be uniformly distributed on the set of 3-connected rooted maps with $n + 4$ edges; then \hat{G}_n converges in distribution to G_∞ in the local weak sense. In particular, G_∞ is almost surely three-connected.

Proof. By Theorem 4.8 and Lemma 5.1 of [6], the conditional distribution of G_n , given that it is 3-connected, is uniform in the set of 3-connected graphs with $n + 4$ edges (note that Q_n has $n + 4$ faces). Furthermore, by Proposition 6.1 of [6], $\mathbf{P}\{G_n \text{ is 3-connected}\} \rightarrow 2^8/3^6$ as $n \rightarrow \infty$. Write $\mathcal{M}(i) = \{M \in \mathcal{M} : M(i) \text{ is irreducible}\}$, and observe that we may then write the event that G_n is 3-connected as $E(M_n) = \{M_n \in \mathcal{M}(U)\}$.


We prove the theorem by showing that for fixed $R > 0$, $E(M_n)$ is asymptotically independent of $M_n^{<R}$. More precisely, we show that any set \mathcal{S} of finite rooted planar maps,

$$|\mathbf{P}\{M_n^{<R} \in \mathcal{S}, E(M_n)\} - \mathbf{P}\{M_n^{<R} \in \mathcal{S}\} \mathbf{P}\{E(M_n)\}| \rightarrow 0, \quad (2)$$

as $n \rightarrow \infty$. Since $M_n^{<R} \rightarrow M_\infty^{<R}$ almost surely, and $G_n^{<R/2}$ is determined by $M_n^{<R}$, the first assertion of the theorem then follows. Having established this, since G_∞ is the distributional limit

of a sequence of random 3-connected maps, it must itself be a.s. 3-connected. It thus remains to prove that $E(M_n)$ and $M_n^{<R}$ are indeed asymptotically independent. For the remainder of the proof, we fix $R > 0$ and a set \mathcal{S} as above.

We again view T_∞ as constructed from the infinite path P and the random variables X_i and random trees B_i . List the nodes of P as $(v_i, i \geq 1)$. Write $V_n^{<R}$ for the set of nodes of T_n that either (a) are nodes of $M_n^{<R}$ or (b) are buds whose attachment corner is incident to a node of $M_n^{<R}$.



(a) Nodes of T_n are grey, edges of P are purple. (b) The trees $T_n(k, 1)$ and $T_n(k, 2)$ (here $k = 1$).

Figure 8. Splitting T_n into $T_n(k, 1)$ and $T_n(k, 2)$.

Given $k > 1$, let $T_n(k, 1)$ be the subtree of T_n containing the root edge when all strict descendants of v_{k+1} are removed, and let $T_n(k, 2)$ be the subtree of T_n consisting of v_k and all its strict descendants; see Figure 8. (If v_{k+1} is not a node of T_n we agree that $T_n(k, 2)$ is empty.) Since T_n is an uniformly random binary tree, conditional on its size $T_n(k, 2)$ is an uniformly random binary tree and is independent of $T_n(k, 1)$.

Let H_n be the set of nodes of T_n incident to a corner κ with $S_{T_n}(\kappa) \leq s^*(T_n) + 7$. For any three-edge path in M_n joining distinct vertices of the hexagon, if the internal nodes of the path do not lie on the hexagon then they are nodes of T_n that neighbour buds which attach to the hexagon. Since buds κ attaching to the hexagon have $S_{T_n}(\kappa) \leq s^*(T_n) + 6$, it follows that all nodes of such a path belong to H_n .

Write $M_n(k, 1)$ and $M_n(k, 2)$ for the closures of $T_n(k, 1)$ and $T_n(k, 2)$. Then conditional on its size, $M_n(k, 2)$ is an uniformly random quadrangulation of the hexagon. Furthermore, by reasoning similar to that in the proof of Corollary 5, it is straightforward to see that for all $\kappa \in \mathcal{C}_B(T_n)$, if κ and $\sigma_{T_n}(\kappa)$ are both elements of $\mathcal{C}_B(T_n(k, 1))$ then $\sigma_{T_n}(\kappa) = \sigma_{T_n(k, 1)}(\kappa)$. Likewise, if $\kappa, \sigma_{T_n}(\kappa) \in \mathcal{C}_B(T_n(k, 2))$ then $\sigma_{T_n}(\kappa) = \sigma_{T_n(k, 2)}(\kappa)$.

Now let $A(k, n)$ be the event that no node of $V_n^{<R}$ is a weak descendant of v_k , that $H_n \subset v(T_n(k, 2))$, and that $|v(T_n(k, 2))| \geq n/2$. Almost surely, $V_n^{<R} = V_\infty^{<R}$ for all n sufficiently large, and $s^*(T_n)$ a.s. decreases to $-\infty$. For any $\epsilon > 0$, we may therefore choose k and n large enough that $\mathbf{P}\{A(k, n)\} > 1 - \epsilon$, and for such k and n we have

$$|\mathbf{P}\{M_n^{<R} \in \mathcal{S}, E(M_n)\} - \mathbf{P}\{M_n^{<R} \in \mathcal{S}, E(M_n), A(k, n)\}| < \epsilon.$$

Furthermore, by the above observations about consistency of closure locations in T_n and $T_n(k, 2)$, if $A(k, n)$ occurs then $M_n^{<R} = M_n^{<R}(k, 1)$ and $E(M_n) = E(M_n(k, 2))$, where we let $E(M_n(k, 2)) = \{M_n(k, 2) \in \mathcal{M}(U)\}$. We thus have

$$\begin{aligned} & \mathbf{P}\{M_n^{<R} \in \mathcal{S}, E(M_n), A(k, n)\} \\ &= \mathbf{P}\{M_n^{<R}(k, 1) \in \mathcal{S}, E(M_n(k, 2)), A(k, n)\} \\ &= \mathbf{P}\{M_n^{<R}(k, 1) \in \mathcal{S}, A(k, n)\} \mathbf{P}\{E(M_n(k, 2)) \mid A(k, n), M_n^{<R}(k, 1) \in \mathcal{S}\} \end{aligned}$$

so

$$\begin{aligned} & |\mathbf{P}\{M_n^{<R} \in \mathcal{S}, E(M_n)\} \\ & - \mathbf{P}\{M_n^{<R}(k, 1) \in \mathcal{S}, A(k, n)\} \cdot \mathbf{P}\{E(M_n(k, 2)) | A(k, n), M_n^{<R}(k, 1) \in \mathcal{S}\}| < \epsilon. \end{aligned}$$

For k and n large enough that $\mathbf{P}\{A(k, n)\} > 1 - \epsilon$, we also have

$$|\mathbf{P}\{M_n^{<R} \in \mathcal{S}\} - \mathbf{P}\{M_n^{<R}(k, 1) \in \mathcal{S}, A(k, n)\}| < \epsilon.$$

Now recall that $T_n(k, 2)$ is an uniform binary tree and is independent of $T_n(k, 1)$ conditional on its size. It follows that given that $|v(T_n(k, 2))| = m$, $M_n(k, 2)$ is distributed as M_m , so

$$\mathbf{P}\{E(M_n(k, 2)) | A(k, n), M_n^{<R}(k, 1) \in \mathcal{S}, |v(T_n(k, 2))| = m\} = \mathbf{P}\{E(M_m)\}.$$

Given $A(k, n)$ we have $n/2 \leq |v(T_n(k, 2))| \leq n$, so by the triangle inequality

$$|\mathbf{P}\{E(M_n(k, 2)) | A(k, n), M_n^{<R}(k, 1) \in \mathcal{S}\} - \mathbf{P}\{E(M_n)\}| \leq \sup_{n/2 \leq m \leq n} 2|\mathbf{P}\{E(M_m)\} - 2^8/3^6|,$$

which is also less than ϵ for n sufficiently large since $\mathbf{P}\{E(M_n)\} = \mathbf{P}\{G_n \text{ is 3-connected}\} \rightarrow 2^8/3^6$ as $n \rightarrow \infty$. The preceding inequalities (and the fact that probabilities lie between zero and one) then yield that for n large,

$$|\mathbf{P}\{M_n^{<R} \in \mathcal{S}, E(M_n)\} - \mathbf{P}\{M_n^{<R} \in \mathcal{S}\} \mathbf{P}\{E(M_n)\}| < 2\epsilon.$$

Since ϵ was arbitrary, this establishes (2). \square

6. Questions and remarks

- (0) It is not hard to show using enumerative results for irreducible quadrangulations with boundary [12] that the degree of the root node has exponential tails in both M_n and G_n .⁴ It then follows, from the general result of Gurel-Gurevich and Nachmias [7] on recurrence of planar graph limits, that simple random walk is recurrent on both graphs.
- (1) A pioneering work of Brooks, Smith, Stone and Tutte [4] showed how to associate to a squaring of a rectangle with any rooted 3-connected map. For random maps, this yields a random squaring. We believe it is possible to show that when appropriately rescaled to remain compact, the squarings corresponding to the sequence G_n , viewed as random subsets of \mathbb{R}^2 converge almost surely for the Hausdorff distance. We are currently pursuing this line of enquiry [1].
- (2) Rather than rooting at e^- , it would also be natural to re-root Q_n at (an orientation of) the edge e^* that is added to the hexagon. However, it is straightforward to see that the sequence (Q_n, e^*) can not converge almost surely. This is because the hexagon essentially corresponds to the corner of minimum label in T_n , and this minimum label tends to $-\infty$ as $n \rightarrow \infty$. Nonetheless, we expect that (Q_n, e^*) does converge in distribution, that the law of the limit can be explicitly described using the results from [6], and that this law is mutually absolutely continuous with that of Q_∞ .
- (3) Marckert [11] essentially establishes that the contour process of T_n and the label process S_{T_n} (in contour order) converge jointly, after appropriate normalization, to a pair (\mathbf{e}, Z) , where \mathbf{e} is a standard Brownian excursion, and Z is the Brownian snake indexed by \mathbf{e} . (See [9] for more details on these objects and their connections with random maps.) Given this, it does not seem out of reach of current technology to prove that Q_n converges, after renormalization, to the Brownian map.

⁴For the assiduous reader: what we call irreducible was called *simple* in [12].

- (4) There is a standard bijection between binary trees and plane trees, that consists of contracting edges from parents to right children in the binary tree; applying this to T_n yields a plane tree P_n . Augmenting the edges of P_n with independent uniform $\{-1, 0, +1\}$ random variables and applying the Schaeffer bijection then yields an uniformly random rooted quadrangulation (not necessarily irreducible) with $n + 1$ nodes. This yields a growth procedure for uniformly random quadrangulations, which I believe deserves investigation.
- (5) Evans, Grübel and Wakolbinger [5] have initiated the study of the Doob-Martin boundary of Luczak and Winkler's tree growth process. Roughly speaking, the Doob-Martin boundary corresponds to the ways in which it is possible to condition on T_∞ to obtain a well-defined conditional growth procedure for $(T_n, n \geq 1)$. It would be interesting to revisit the growth of M_n and G_n in this context.

7. Acknowledgements

My sincere thanks to Nicholas Leavitt, for stimulating conversations and for his helpful comments on a draft of this work. Thanks also to Marie Albenque and Gilles Schaeffer, who were the first to teach me about the bijection between T_n and M_n , in 2011.

References

- [1] Louigi Addario-Berry and Nicholas Leavitt. A random infinite squaring of a rectangle. In preparation, 2014+.
- [2] David Aldous and J. Michael Steele. The objective method: probabilistic combinatorial optimization and local weak convergence. In *Probability on discrete structures*, volume 110 of *Encyclopaedia Math. Sci.*, pages 1--72. Springer, 2004. URL <http://www.stat.berkeley.edu/~aldous/Papers/me101.pdf>.
- [3] Itai Benjamini and Oded Schramm. Recurrence of distributional limits of finite planar graphs. *Electron. J. Probab.*, 6:no. 23, 13 pp. (electronic), 2001. URL <http://arxiv.org/abs/math/0011019>.
- [4] R. L. Brooks, C. A. B. Smith, A. H. Stone, and W. T. Tutte. The dissection of rectangles into squares. *Duke Math. J.*, 7:312--340, 1940.
- [5] Steven N. Evans, Rudolf Grübel, and Anton Wakolbinger. Trickle-down processes and their boundaries. *Electron. J. Probab.*, 17:no. 1, 58, 2012. URL <http://arxiv.org/abs/1010.0453>.
- [6] Éric Fusy, Dominique Poulalhon, and Gilles Schaeffer. Dissections, orientations, and trees with applications to optimal mesh encoding and random sampling. *ACM Trans. Algorithms*, 4(2):19:1--19:48, 2008. URL <http://arxiv.org/abs/0810.2608>.
- [7] Ori Gurel-Gurevich and Asaf Nachmias. Recurrence of planar graph limits. *Ann. of Math. (2)*, 177(2): 761--781, 2013. URL <http://arxiv.org/abs/1206.0707>.
- [8] Sergei K. Lando and Alexander K. Zvonkin. *Graphs on surfaces and their applications*, volume 141 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2004. ISBN 3-540-00203-0.
- [9] Jean-François Le Gall and Grégory Miermont. Scaling limits of random trees and planar maps. In *Probability and statistical physics in two and more dimensions*, volume 15 of *Clay Math. Proc.*, pages 155--211. Amer. Math. Soc., 2012. URL <http://www.math.u-psud.fr/~jfllegall/Cours-Buziosf.pdf>.
- [10] Malwina Luczak and Peter Winkler. Building uniformly random subtrees. *Random Structures Algorithms*, 24(4):420--443, 2004. URL <http://www.math.dartmouth.edu/~pw/papers/birds.ps>.
- [11] Jean-François Marckert. The rotation correspondence is asymptotically a dilatation. *Random Structures Algorithms*, 24(2):118--132, 2004. URL <http://www.labri.fr/perso/marckert/propre.pdf>.
- [12] R. C. Mullin and P. J. Schellenberg. The enumeration of c -nets via quadrangulations. *J. Combinatorial Theory*, 4:259--276, 1968. URL <http://www.sciencedirect.com/science/article/pii/S0021980068800079#>.

Department of Mathematics and Statistics, McGill University, 805 Sherbrooke Street West, Montréal, Québec, H3A 2K6, Canada

E-mail address: louigi@math.mcgill.ca

URL: <http://www.math.mcgill.ca/~louigi/>